



REVYPEEL® HIGH & LOW

MEDICAL PROTOCOLS



This document contains the official in-clinic application protocols for REVYPEEL® LOW and REVYPEEL® HIGH. It is intended for dermatologists, aesthetic physicians, and other qualified medical professionals.

General Warnings

This document contains the official in-clinic application protocols for REVYPEEL® LOW and REVYPEEL® HIGH. It is intended for dermatologists, aesthetic physicians, and other qualified medical professionals.



To preserve the chemical and physical properties of the product, do not open the bottle. Extract REVYPEEL® using a disposable syringe by piercing the grey rubber cap with the rounded-tip applicator provided.



At the end of the treatment, apply an SPF 50+ cream or a total sunblock to prevent the onset of hyperpigmentation.



The treatment can be safely performed even during the summer without damaging the tissue. In warmer months, it is advisable to store the product in a refrigerator. Use at room temperature.



Use nitrile gloves during application.



Protect mucous membranes, perinasal area, eye contour, tattoos, cuts, other skin lesions, and any suspicious lesions with a barrier cream or petroleum jelly. (ADD to the LIST of general Advice and Warnings)



Start application on a limited area. A slight tingling sensation may occur. If the patient experiences a strong burning sensation, stop the treatment, neutralize with REVYSKIN Neutralizing Gel, rinse the treated area with cold water, and apply REVYSKIN Post Peeling Cream SPF 50+.



Avoid the periocular area to prevent potential irritation.



Perform the treatment only on intact skin and at a reasonable distance from hair removal, shaving, or other forms of stimulation.



Avoid treating areas affected by seborrheic dermatitis.



Avoid the corners of the mouth (especially in the presence of angular cheilitis) and the area around the nose.



Use caution when treating the inframammary fold due to potential fissures in macerated skin.



On acne-prone or oily skin and on delicate facial areas (e.g., sides of the mouth), limit the application to a maximum of three passes during the first session. Massage vigorously to promote effective penetration.



NOTE: The protocols provided below serve as general guidelines. They can be customized by the practitioner and used in synergy with other techniques such as micro or nano needling, PRP, skin boosters, carboxytherapy, PICO etc. Please refer to the Combined protocols in the second section of this guideline.

REVYPEEL® LOW

Protocol for:

TREATMENT OF FACIAL, NECK, AND DÉCOLLETÉ REVITALIZATION INFLAMMATORY ACNE

Protocol Summary:



INDICATIONS

Facial

Neck

Décolleté Revitalization

Inflammatory Acne



TREATMENT FREQUENCY

1 session/week
for 5 weeks



QUANTITY TO USE

2 ml



FITZPATRICKS SKIN TYPE

From I to VI



CONTACT TIME

5 - 8 minutes

APPLICATION PROTOCOL

- 1 Wear nitrile gloves. Cleanse the skin thoroughly and remove any makeup.
- 2 Protect mucous membranes, perinasal area, eye contour, tattoos, cuts, other skin lesions, and any suspicious lesions with a barrier cream or petroleum jelly.
- 3 Prepare the skin by soaking a cotton pad with **REVYSKIN PRE PEELING LOTION** and applying it evenly to the desired areas.
- 4 Withdraw 2 ml of **REVYPEEL® LOW** by piercing the grey rubber cap using either a sterile needle or the provided rounded applicator tip. Apply the solution zone by zone to the treatment area.
- 5 Apply a few drops of **REVYPEEL® LOW** to each area and massage with firm friction.
- 6 Repeat the application until visible plumpness and turgidity is observed. On average, three passes are needed for a noticeable result: fewer passes (1–2) are recommended for the neck and very sensitive or delicate skin; more passes (up to 5) may be required for thicker skin.
- 7 Continue the application zone by zone until the entire surface has been treated, avoiding the eyelid area.
- 8 Leave the product on for **5 to 8 minutes** per area approximately.
- 9 Begin neutralizing the peel by gently applying water with a cotton pad using circular motions. Follow immediately with **REVYSKIN NEUTRALIZING GEL**, blending it in with circular movements to activate the neutralization process. Allow the product to act for a few minutes, then rinse thoroughly with cold water.
- 10 Apply **REVYSKIN POST PEELING CREAM 50+** to the treated areas.
- 11 Advise the patient to continue at-home care with **REVYSKIN POST PEELING CREAM SPF 50+**.



NOTE: In the treatment of **active acne**, if some acne scars remain after completing the full treatment cycle, it is recommended to perform a few additional sessions with **REVYPEEL® HIGH**, targeting the affected scarred areas.

ROSACEA: Rosacea should always be approached as a highly sensitive skin condition.

REVYPEEL® LOW can be a valuable option for managing this condition, including papulopustular rosacea.

However, it is strongly recommended not to leave the peel on the affected area for more than 5 minutes.

REVYPEEL® HIGH

Protocol for:

INTENSE FACE, NECK & DÉCOLLETÉ REVITALIZATION

STRETCH MARKS

ATROPHIC SCARS (ACNE, CHICKENPOX)

Protocol Summary:



INDICATIONS

Intense Face

Neck

Décolleté Revitalization

Stretch-Marks

Atrophic Scars (acne, chickenpox)



TREATMENT FREQUENCY

Every 14 days for 5 sessions and/or until lesion improvement is achieved



QUANTITY TO USE

2 ml



FITZPATRICKS SKIN TYPE

From I to III



CONTACT TIME

3 - 5 minutes



NOTE: The treatment of scars is suitable in combination with fractional lasers, microneedling, Oxyneedling, and Carboxytherapy).



WARNING: Use REVYPEEL® HIGH with caution on Fitzpatrick skin type IV. Minimize the number of passes and reduce exposure time considering prior skin preparation with tyrosinase inhibitors to limit the risk of adverse reactions.

APPLICATION PROTOCOL

- 1 Wear nitrile gloves. Cleanse the skin thoroughly and remove any makeup.
- 2 Protect mucous membranes, perinasal area, eye contour, tattoos, cuts, other skin lesions, and any suspicious lesions with a barrier cream or petroleum jelly.
- 3 Prepare the skin by soaking a cotton pad with **REVYSKIN PRE PEELING LOTION** and applying it evenly to the desired areas.
- 4 Withdraw 2 ml of **REVYPEEL® HIGH** by piercing the grey rubber cap using either a sterile needle or the provided rounded applicator tip. Apply the solution zone by zone to the treatment area.
- 5 Apply a few drops (1 to 3) of **REVYPEEL® HIGH** to each area and massage with firm friction.
- 6 Repeat the application until visible plumpness and turgidity is observed. On average, three passes are needed for a noticeable result: fewer passes (1–2) are recommended for the neck and very sensitive or delicate skin; more passes (up to 5) may be required for thicker skin.
- 7 Continue the application zone by zone until the entire surface has been treated, avoiding the eyelid area.
- 8 Leave the product on for **3 to 5 minutes** per area approximately.
- 9 Begin neutralizing the peel by gently applying water with a cotton pad using circular motions. Follow immediately with **REVYSKIN NEUTRALIZING GEL**, blending it in with circular movements to activate the neutralization process. Allow the product to act for a few minutes, then rinse thoroughly with cold water.
- 10 Apply **REVYSKIN POST PEELING CREAM 50+** to the treated areas.
- 11 Advise the patient to continue at-home care with **REVYSKIN POST PEELING CREAM SPF 50+**, or an alternative broad-spectrum sunscreen.

REVYPEEL® LOW | REVYPEEL® HIGH

Protocol for:

SKIN HYPERPIGMENTATION AND DYSCHROMIA

Protocol Summary:



INDICATIONS

Skin Hyperpigmentation

Dyschromia



TREATMENT FREQUENCY

1 application every
4 weeks



QUANTITY TO USE

2 ml



FITZPATRICKS SKIN TYPE

REVYPEEL® LOW: IV to V

REVYPEEL® HIGH: I to III



CONTACT TIME

3 - 5 minutes



WHEN TO APPLY THE PEEL

Nanoneedling: before

Microneedling,
Oxyneedling: after



NOTE: The use of REVYPEEL® HIGH is recommended for Fitzpatrick skin types I to II and the use of REVYPEEL® LOW is recommended for Fitzpatrick skin types IV to VI


APPLICATION PROTOCOL

- 1 Wear nitrile gloves. Cleanse the skin thoroughly and remove any makeup.
- 2 Protect mucous membranes, perinasal area, eye contour, tattoos, cuts, other skin lesions, and any suspicious lesions with a barrier cream or petroleum jelly.
- 3 Apply a lidocaine-based topical anaesthetics to the treatment area. Remove it after the recommended exposure time.
- 4 Prepare the skin by soaking a cotton pad with REVYSKIN PRE PEELING LOTION and applying it evenly to the desired areas.
- 5 Withdraw 2 ml of REVYPEEL® LOW or REVYPEEL® HIGH, depending on the skin phototype, by piercing the grey rubber cap using either a sterile needle or the provided rounded applicator tip. Apply the solution zone by zone to the treatment area.
- 6 Repeat the application until visible plumpness and turgidity is observed. On average, three passes are needed for a noticeable result: fewer passes (1–2) are recommended for the neck and very sensitive or delicate skin; more passes (up to 5) may be required for thicker skin.
- 7 Leave the product on for 3 to 5 minutes per area approximately.
- 8 Begin neutralizing the peel by gently applying water with a cotton pad using circular motions. Follow immediately with REVYSKIN NEUTRALIZING GEL, blending it in with circular movements to activate the neutralization process. Allow the product to act for a few minutes, then rinse thoroughly with cold water.
- 9 Perform a **Micro Needling, Oxyneedling** (using needles no longer than 0.25 mm), or Nanoneedling treatment on the targeted area following standard techniques.
- 10 Apply REVYSKIN POST PEELING CREAM 50+ to the treated areas.
- 11 Advise the patient to continue at-home care with REVYSKIN POST PEELING CREAM SPF 50+, or an alternative broad-spectrum sunscreen.



NOTE: In the event of side effects causing persistent inflammation (erythema lasting more than 24 hours), especially in predisposed individuals, there is a risk of post-inflammatory hyperpigmentation. Ensure they avoid sun exposure, instruct them to use a high-protection sunscreen, and consider prescribing tyrosinase inhibitors or topical corticosteroids.

GENERAL CONTRAINDICATIONS FOR THE APPLICATION OF REVYPEEL® LOW AND REVYPEEL® HIGH




Melasma



Hypertrophic or
keloid scars



Broken skin



Known allergy
to any of the
product's
ingredients



Pregnancy and
breastfeeding



Active skin rashes
and/or herpetic
outbreaks

Potential Side Effects and Adverse Reactions

- **Intense exfoliation:** This may occur during the initial applications, particularly on skin with a thick stratum corneum. It can last up to 10 days. Apply **REVYSKIN POST PEELING CREAM** to restore, hydrate, and rebalance the treated skin.
- **Severe burning sensation immediately after application:** This may occur in individuals with very sensitive skin. Rinse the area immediately with cool running water and apply **REVYSKIN POST PEELING CREAM**.
- **Immediate erythema:** Typically resolves within a few hours. Application of **REVYSKIN POST PEELING CREAM** is **recommended**.
- **Excessive dryness:** Apply **REVYSKIN POST PEELING CREAM** to hydrate, rebalance, and nourish the treated tissue.
- **Contact with the conjunctiva:** Rinse with sterile saline solution.
- **Itching, with or without erythema, occurring several days after treatment** is a very rare event and may indicate an allergic reaction to kojic acid. In such cases, consider prescribing a topical corticosteroid.
- **Delayed localized inflammation (appearing the following day):** Presents as redness and/or swelling in a limited area of treated skin and is likely due to irritation on skin that was not fully intact at the time of application. Prescribe a topical corticosteroid.
- **Appearance of darker patches of skin in the days following treatment:** These are caused by desiccation of thicker skin areas and typically resolve spontaneously with desquamation.



NOTE: In the event of **side effects causing persistent inflammation** (erythema lasting more than 24 hours), especially in predisposed individuals, there is a risk of **post-inflammatory hyperpigmentation**. Ensure they avoid sun exposure, instruct them to use a high-protection sunscreen, and consider prescribing **tyrosinase inhibitors** or **topical corticosteroids**.



REVYPEEL® HIGH & LOW

PROTOCOLS COMBINED
WITH OTHER TECHNIQUES

This document contains the official in-clinic application protocols for REVYPEEL® LOW and REVYPEEL® HIGH. It is intended for dermatologists, aesthetic physicians, and other qualified medical professionals.

PROTOCOLS COMBINED WITH OTHER TECHNIQUES

Nanoneedling,
Microneedling,
Oxyneedling

Cryotherapy

Polydioxanone
(PDO) Threads

Laser
therapies

PRP
(Platelet-Rich
Plasma)

Fractional
Laser

Carboxytherapy

IPL

Injectable
Biorevitalization

REVYPEEL® LOW NANO NEEDLING

Protocol Summary:



INDICATIONS

Actinic Keratoses

Age Spots

Post-Acne
Hyperpigmentation



TREATMENT FREQUENCY

1 session/week for
5 weeks



QUANTITY TO USE

1 -1,5 ml



FITZPATRICKS SKIN TYPE

From I to VI



CONTACT TIME

5 minutes



WHEN TO APPLY THE PEEL

After

APPLICATION PROTOCOL

- 1 Wear nitrile gloves.
- 2 Cleanse the skin.
- 3 Perform the nano needling treatment by executing circular motions on and around the target spot until mild erythema appears.
- 4 Soak a cotton pad with **REVYSKIN PRE PEELING LOTION** and apply it to the treatment area.
- 5 Draw **1 ml** from the **REVYPEEL® LOW** vial by piercing the grey rubber stopper with the rounded applicator provided. Apply 1 drop of product and manually massage until completely absorbed. Repeat the application for a total of 3 passes. Leave on for **5 minutes**.
- 6 Neutralize the peeling by applying **REVYSKIN NEUTRALIZING GEL** with a brush. Use circular motions to blend the two products and activate neutralization. Leave on for a few minutes, then rinse with cold water.
- 7 Apply **REVYSKIN POST PEELING CREAM SPF 50+** to the treated areas.
- 8 Recommend continued home use of **REVYSKIN POST PEELING CREAM SPF 50+** or another broad-spectrum, high-protection sunscreen.



MICRONEEDLING and OXY NEEDLING

Protocol Summary:



INDICATIONS

Atrophic Scars from Acne and Chickenpox

Scars

Stretch Marks



TREATMENT FREQUENCY

1 session every 10–15 days (on repaired and intact skin) for 5–8 sessions



QUANTITY TO USE

1 - 1,5 ml



FITZPATRICKS SKIN TYPE

From I to VI



CONTACT TIME

2 - 5 minutes



WHEN TO APPLY THE PEEL

Before

APPLICATION PROTOCOL

- 1 Wear nitrile gloves.
- 2 Cleanse the skin.
- 3 Apply a lidocaine-based topical anaesthetic to the treatment area. Remove it after the appropriate exposure time.
- 4 Soak a cotton pad with **REVYSKIN PRE PEELING LOTION** and apply it to the treatment area.
- 5 Depending on the size of the area to be treated, draw **1–1.5 ml** from the **REVYPEEL® HIGH** vial by piercing the grey rubber stopper with the rounded applicator provided. Apply 2–3 drops and massage manually until fully absorbed. Repeat the application for a total of 3 passes. Leave on for 2–5 minutes.
- 6 Neutralize the peeling by applying **REVYSKIN NEUTRALIZING GEL** with a brush, using circular movements to blend and activate the neutralization. Leave on for a few minutes, then rinse with cold water.



WARNING: In particularly sensitive or reactive patients, if there is discomfort upon contact between Neutralizing Gel and REVYPEEL® HIGH, first remove REVYPEEL® HIGH with a cloth soaked in cold water at the end of the exposure time. Then apply the Neutralizing Gel to the skin, massage vigorously for one minute, and remove with a cloth soaked in cold water.

- 7 Proceed with the **microneedling or Oxyneedling treatment**, performing circular and linear movements over and around the scar.
- 8 Cleanse the treated area and apply gauze soaked in cold water. If needed, apply ice packs.
- 9 Apply a specific healing cream.
- 10 Starting the day after treatment, apply **REVYSKIN POST PEELING CREAM SPF50+**.
- 11 Recommend continued home use of **REVYSKIN POST PEELING CREAM SPF50+**.



PDO THREADS LIFTING

Protocol Summary:



INDICATIONS

Threads lifting



QUANTITY TO USE

1,5 ml



FITZPATRICKS SKIN TYPE

From I to VI



CONTACT TIME

REVYPEEL® LOW: 8 min

REVYPEEL® HIGH: 5 min



WHEN TO APPLY THE PEEL

Before

APPLICATION PROTOCOL

- 1 Cleanse the skin thoroughly.
- 2 Apply a lidocaine-based topical anesthetic to the area to be treated. Remove it after the appropriate exposure time.
- 3 Prepare the skin by soaking a cotton pad with **REVYSKIN PRE PEELING LOTION** and applying it evenly to the treatment area.
- 4 Draw **1,5 ml** of **REVYPEEL® LOW** or **REVYPEEL® HIGH** by piercing the grey rubber stopper with the rounded applicator tip provided. Proceed with the treatment zone by zone.
- 5 Apply a few drops of product to each area and massage vigorously with firm friction.
- 6 Repeat the application for a total of **3 passes**. Leave on for **8 minutes** (for **REVYPEEL® LOW**) or **5 minutes** (for **REVYPEEL® HIGH**).
- 7 Neutralize the peel by applying **REVYSKIN NEUTRALIZING GEL** using a brush and performing circular motions to blend the two products and enhance the neutralizing effect. Allow a few minutes to act, then rinse with cold water.
- 8 Apply **REVYSKIN POST PEELING CREAM SPF 50+** to the treated areas.
- 9 Proceed with the implantation of PDO (polydioxanone) threads.
- 10 Upon completion of the procedure, apply ice packs to the treated areas for a few minutes to reduce inflammation and soothe the skin.

REVYPEEL® HIGH

PRP (PLATELET-RICH PLASMA)

Protocol Summary:



INDICATIONS

Alopecia



QUANTITY TO USE

0,5 - 1 ml



FITZPATRICKS SKIN TYPE

From I to VI



CONTACT TIME

10 minutes



WHEN TO APPLY THE PEEL

Before - Perform the PRP the following day

APPLICATION PROTOCOL

- 1 Apply a few drops (0.5–1 ml) of **REVYPEEL® HIGH** uniformly to the scalp over the alopecic areas, parting the hair to ensure direct contact with the skin.
- 2 Perform **a single pass** and massage the product into the scalp until fully absorbed.
- 3 Leave on for **10 minutes**.
- 4 Rinse with cold water.
- 5 On the following day, perform the **injectable PRP (Platelet-Rich Plasma)** treatment.



REVYPEEL®

REVYPEEL® LOW | REVYPEEL® HIGH

CARBOXYTHERAPY

Treatment with REVYPEEL® LOW or REVYPEEL® HIGH may be performed prior to subcutaneous carboxytherapy within the same session, to enhance overall skin quality and its immediate response. This combined approach results in a visible improvement in skin radiance and turgor.

The subsequent carboxytherapy application, by enhancing local blood circulation and stimulating fibroblast activity, further improves skin elasticity and contributes to tissue toning and firming.

Protocol Summary:



INDICATIONS

All the indications for carboxytherapy



QUANTITY TO USE

Face 1,5 ml
Stretch-marks or body
3,5 ml



CONTACT TIME

5 - 10 minutes



WHEN TO APPLY THE PEEL

Before, in the same session

APPLICATION PROTOCOL

- 1 Perform the complete **REVYPEEL® LOW or REVYPEEL® HIGH** treatment, depending on the area and condition being treated, following the previously described protocol.
- 2 Proceed with the **carboxytherapy session**, in accordance with the most appropriate clinical protocol.



REVYPEEL® LOW | REVYPEEL® HIGH

INJECTABLE, BIOREVITALIZATION or BOTOX

Protocol Summary:



INDICATIONS

Filler, biorevitalization or botox



TREATMENT FREQUENCY

1 session per month for a total of 5 session



QUANTITY TO USE

1 ml



FITZPATRICKS SKIN TYPE

From I to VI



CONTACT TIME

REVYPEEL® LOW: 5-8 min

REVYPEEL® HIGH: 5 min



WHEN TO APPLY THE PEEL

INJECTABLE BIOREVITALISATION:
Before, same session

BOTOX: Before at least **one week** prior to Botox

APPLICATION PROTOCOL

- 1 Cleanse the skin thoroughly.
- 2 Prepare the tissue by soaking a cotton pad with **REVYSKIN PRE PEELING LOTION** and applying it evenly over the treatment area.
- 3 Draw 1 ml of **REVYPEEL® LOW** or **REVYPEEL® HIGH** from the vial by piercing the grey rubber stopper with the rounded applicator tip provided, and proceed with treatment zone by zone.
- 4 Apply a few drops of the product to each area and massage with firm friction.
- 5 Repeat the application for a total of 3 passes. Leave on for **5–8 minutes (REVYPEEL® LOW)** or **5 minutes (REVYPEEL® HIGH)**.
- 6 Neutralize the peeling by applying **REVYSKIN NEUTRALIZING GEL** with a brush. Use circular movements to blend the two products and activate the neutralization process. Allow it to act for a few minutes, then rinse with cold water.
- 7 Apply **REVYSKIN POST PEELING CREAM SPF 50+** to the treated areas.
- 8 Proceed with the **injectable biorevitalization** treatment by administering micro-papular intradermal injections in the same area previously treated with REVYPEEL®.



NOTE: Botulinum toxin injections (BOTOX) should be performed in a separate session, at least one week after the REVYPEEL® treatment.

REVYPEEL® LOW

IPL (Intense Pulsed Light)

Protocol Summary:



CONTACT TIME

5 minutes



WHEN TO APPLY THE PEEL

After the IPL session

APPLICATION PROTOCOL

- 1 Perform the **IPL session** in accordance with the most appropriate clinical protocol.
- 2 Proceed the complete **REVYPEEL® LOW** treatment, depending on the area and condition being treated, following the previously described protocol.
- 3 **Contact time:** 5 minutes



NOTE: REVYPEEL® LOW is generally well tolerated by **Asian skin types**.

However, in cases where the practitioner deems it appropriate, patients with **Fitzpatrick phototypes IV to VI** may undergo **pre-treatment with tyrosinase inhibitors** during the two months preceding the peeling procedure.



REVYPEEL® LOW

FRACTIONAL RADIOFREQUENCY

Protocol Summary:



INDICATIONS

Treatment of scarring or to be performed at every RF microneedling session



CONTACT TIME

5 minutes



WHEN TO APPLY THE PEEL

Before, same session

APPLICATION PROTOCOL

- 1 Proceed the complete **REVYPEEL® LOW** treatment, depending on the area and condition being treated, following the previously described protocol.
- 2 **Application time:** 5 minutes
- 3 Apply a lidocaine-based topical anaesthetic to the area to be treated. Remove it after the appropriate exposure time.
- 4 Perform the **FRACTIONAL RF** session in accordance with the most appropriate clinical protocol.



REVYPEEL® LOW LASER THERAPIES

REVYPEEL® LOW protocol can be perfectly combined **AFTER** several laser therapies and for the following indications:

LASER	INDICATION	REVYPEEL® LOW SESSION (before or after)
IPL	ROSACEA	AFTER
PICO LASER	PIGMENTATION/ REJUVENATION	AFTER
CO ₂ LASER	SCARRING (NOTE)	NOT RECOMMENDED



NOTE: For scarring we would recommend using REVYPEEL® with **microneedling** in those **patients who don't want lasers** or those who have **darker skin** and are contraindicated to this laser.

Protocol Summary:



INDICATIONS

IPL: Rosacea

PICO LASER:
Pigmentation /
Rejuvenation

CO₂ LASERS: Scarring



CONTACT TIME

5 minutes



WHEN TO APPLY THE PEEL

After
CO₂ LASER is **not recommended** in combination with Revypeel®

APPLICATION PROTOCOL

- 1 Perform the **LASER** session in accordance with the most appropriate clinical protocol.
- 2 Proceed the complete **REVYPEEL® LOW** treatment right after lasers in the same session, depending on the area and condition being treated, following the previously described protocol.
- 3 **Application time:** 5 minutes



WARNING: Fitzpatrick Skin Types IV–VI: For darker skin phototypes (IV to VI) we would recommend pre-treating the skin for 2–3 months with agents such as hydroquinone or arbutin to minimize the risk of post-treatment pigmentation or inflammation. Then use **REVYPEEL® LOW** protocol.

REVYPEEL® HIGH CRYOTHERAPY

Protocol Summary:



INDICATIONS

Solar Lentigo

Actinic Keratosis



WHEN TO APPLY THE PEEL

Before

APPLICATION PROTOCOL

- 1 Perform a complete **REVYPEEL® HIGH** treatment, depending on the area and the condition being addressed, following the previously outlined protocol.
- 2 Proceed with the **cryotherapy** session, in accordance with the most appropriate clinical protocol.

